

**Helping your child live a  
healthy life with asthma**



**Managing  
your child's  
asthma**

**Asthma  
+ Respiratory**  
FOUNDATION NZ

**The hongī symbolises  
the breath of life.**

**It recalls the moment  
when Tāne breathed life  
into Hīneahuone,  
the first earthly woman.**

---

# Introduction

Asthma is a common New Zealand illness, but it's scary when it affects the child you care for. The good news is that asthma can be well managed.

This booklet is for the parents, whānau, and other caregivers of children with asthma. It will help you make sure your child stays fit, healthy, and happy. You will learn how to help prevent an asthma attack and what to do when one takes place.

There's also a digital resource online at:

**[www.learnaboutlungs.org.nz](http://www.learnaboutlungs.org.nz)**

It includes video clips and animations that you and your child could view together.

You don't need to read the booklet all at once. You can just dip in and out of it. It will help you to answer these questions:


■ What is asthma?	2
■ What are the symptoms of asthma?	5
■ What are the triggers for asthma?	7
■ How is asthma treated?	11
■ What should I do to manage my child's asthma?	21
■ What should I do in an emergency?	23
■ Should my child be physically active?	25
■ Who can help?	27
■ Who are our asthma health providers?	33

---

## What is asthma?

Asthma is a common illness of the respiratory system. In New Zealand, one child in seven needs to take medicine for asthma.

### The respiratory system



The respiratory system is the system through which we breathe. We breathe in to bring oxygen into our bodies, and we breathe out to remove carbon dioxide. The respiratory system includes the nose, mouth, lungs, and a set of airways.


The largest airway is the windpipe, which connects the lungs to the nose and mouth. There are more airways inside the lungs. The airways are also called breathing tubes.

People with asthma have sensitive airways. Your child's sensitive airways may be irritated by a trigger in the environment (such as pollen), by an illness, or by an emotion (such as anxiety). The irritation makes your child's airways swell up so that it becomes difficult to breathe. You can read more about asthma triggers on pages 7–10.


## What happens in an asthma attack?

Most people can keep their asthma under control for most of the time. You can teach your child to notice the signs of asthma and what to do to stop having a full asthma attack or 'flare-up'.

Knowledge is power and you can use this power to help prevent your child having an asthma attack. You need to know your child's body, know what triggers your child's asthma, and make sure your child always takes their medicine.



In this **healthy airway**, the tube is wide open so air can easily pass through.



This is the **airway of a person who is having an asthma attack**. The muscles around the airway have tightened up. The lining is swollen and the tube has filled with thick, sticky mucus. All this makes it hard for the person to breathe.



Renee's son Alex has a severe form of asthma that's complicated by a restriction of the throat, but he's also a keen sportsman.



Alex's asthma is serious, but it's never stopped him pursuing his passion for sport! There's been running, football, and underwater hockey, and these days he's into rowing.

I've been so impressed with how Alex has taken it all in his stride. Even when he was quite young, he's wanted to work with us all to find out how to stop getting sick and get things right again when he is. These days, he confidently discusses his action plan with his doctor and practice nurse and takes on a lot of the responsibility for carrying them out.



<sup>1</sup> You can hear more from Renee by going to the digital resource located at [www.learnaboutlungs.org.nz](http://www.learnaboutlungs.org.nz)

---

# What are the symptoms of asthma?

It's important to know the symptoms of asthma. Asthma can usually be quickly managed, but it is dangerous if it is just left.

Asthma has four main symptoms:

- coughing
- breathlessness
- wheezing
- a tight feeling in the chest



**Wheezing** is a whistling sound that people make when they are trying to breathe through airways that are narrower than usual. It is often louder when trying to breathe out.

In an asthma attack, these symptoms get worse, so the person finds it difficult to breathe. Some people say it feels like they have something heavy on their chest or it feels tight. It can be so hard to breathe that it becomes painful.

## What does it feel like?



To get an idea of how asthma feels for your child, try breathing in and out through a straw. First do it normally. Then pinch the straw's end and keep trying to breathe in and out. You will start to feel short of breath and you may feel tightness in your chest.

If you don't have a straw, try blowing air through an open fist. Then gradually close your fist, still trying to blow through it. Imagine how tiring this would feel!

Very young children can't tell people how they're feeling. They need other people to keep an eye out and notice when symptoms start to appear.

You might notice that your child is getting tired because their difficulty in breathing makes it hard to sleep. In a more severe case, you might see your child's chest and sides pulling in.

With an older child, you can ask how they're feeling and help them to watch out for asthma symptoms for themselves.

Your doctor may give you a symptom diary. You and the doctor can use this diary to keep track of what is happening and understand the patterns in your child's symptoms. You can read more about symptom diaries on page 21.



---

## What are the triggers for asthma?

In the past, some people thought that asthma was ‘all in the head’. Now we know that it is the body’s response to a trigger that causes someone’s airways to become narrow.

The triggers can be different for different people, but there are some that are very common.

In the left-hand side of the table on pages 8, 9, and 10, you can see some common triggers for asthma. In the right-hand column, there are some suggestions for how you can help your child avoid these triggers.

If you wish, you can use the middle column to note the possible triggers for your child’s asthma. Mark the column with a tick for ‘yes’, a cross for ‘no’, and a question mark if you’re not sure.

Talk to other people who know your child well about possible triggers they’ve noticed.



Some of the asthma triggers, like pollen and pet fur, contain allergens. Allergens are tiny substances that are usually harmless, but they can cause an allergic reaction in some people.

<b>Common asthma triggers</b>	<b>Your child's triggers</b>	<b>Some ways to help your child avoid asthma triggers</b>
	<input type="checkbox"/> <input checked="" type="checkbox"/> <input type="checkbox"/>	
<b>Colds and flu</b> 	<input type="checkbox"/>	<p>All children get colds. When this happens, watch carefully for signs of asthma. Be ready to change your child's medicine in line with their action plan (page 21).</p> <p>Make sure your child is immunised in autumn each year. The flu vaccination is free to all children who are on preventative asthma medication.</p>
<b>Cigarette smoke</b> 	<input type="checkbox"/>	<p>Keep your child's environment auahi kore (smokefree). There's lots of help if you want to give up smoking. Start with your doctor or nurse or try Quitline. (See page 35 for contact details.)</p>
<b>Cold dry air or changes in the weather</b> 	<input type="checkbox"/>	<p>Keep your child warm and dry and use a scarf to cover their mouth and nose. Encourage your child to warm the air by breathing in through their nose instead of their mouth. If their nose is usually blocked, discuss this with your doctor.</p>
<b>Stress and anxiety</b> 	<input type="checkbox"/>	<p>Do what you're already doing to keep your child strong and healthy through proper rest, good food, exercise, and taking the time to talk through any worries. Relaxed breathing reduces stress and anxiety.</p>
<b>Mould</b> 	<input type="checkbox"/>	<p>Try to keep your house warm, dry, and well aired. Vent your laundry dryer to the outside and use extractor fans in the kitchen and bathroom. If possible, use a dehumidifier. Treat any mould with diluted household bleach, but be careful not to breathe in the fumes.</p>

## Common asthma triggers

## Your child's triggers

## Some ways to help your child avoid asthma triggers



### The pollen in some plants



Pollen is a common trigger, especially in summer and in the early morning and evening. Try to keep outdoor activities for the early afternoon. Close windows when it is windy or humid. Choose pretty, brightly coloured plants for the garden that use insects to transfer their pollen, instead of the wind.

### Cats and other furry pets



Don't get a pet without first checking whether your child is allergic to it. If you can't avoid having a furry pet, talk to your vet about how to keep it well groomed. If you go to a house where there is a furry pet, ask for the pet to be put outside. Your child should never sleep in a room with a pet.

If you move into a house that has had a pet, you will need to clean it thoroughly. It can take three to four months of regular deep cleaning to get rid of all the allergens.

### Dust mites



All houses have some dust mites, but there are ways you can keep them down. For example, you can:

- vacuum each week using a vacuum cleaner with a good filter system
- wash bed linen weekly in hot water
- wash soft toys in hot water or freeze them for 72 hours
- air bed linen and rugs in the sun
- keep your house light and well aired
- dust with a damp cloth

### Physical activity



Exercise is good for everybody, including people with asthma! However, it can trigger asthma, especially when the air is cold and dry. On page 25 there is information about how your child can exercise safely and happily.

## An unusual asthma trigger

## Your child's triggers

## Some ways to help your child avoid this trigger



### Food



Very few children are allergic to food like cow's milk, wheat, seafood, eggs, soy, and peanuts. If you think a food is triggering your child's asthma, visit your health provider (for examples of health providers, see pages 33–35). There are simple tests the health provider can run to check whether food is the problem. If it is, your health provider can suggest other healthy foods to replace it.

You know your child better than anyone. You can use your knowledge to help your child avoid the triggers for their asthma. Your child won't always be able to avoid their triggers, but by knowing what they are, you can help manage them.



Most kids will be exposed to triggers like exercise, animals, and grass, and sometimes that will mean their asthma plays up. I don't think that parents need to feel guilty that their child with asthma is living a normal life.



Tāne, asthma expert

---

# How is asthma treated?


Asthma cannot be cured, but it can be treated. In this part of the booklet, you will read about asthma medicine and the way it is delivered.

## Introducing inhalers

### What is an inhaler?

An asthma **inhaler** is a plastic case that holds asthma medicine. It is designed to deliver the medicine directly to the lungs. Inhalers are often called **puffers**.

There are two types of inhaler: metered dose inhalers (MDIs) and dry powder inhalers.



### Metered dose inhaler

**A metered dose inhaler (MDI)** delivers a set amount of asthma medicine. The MDI contains a pressurised canister that holds the medicine in the form of gas. The MDI needs to be shaken to mix the medicine before it is used. When the MDI is pressed, the canister releases its gas into a spacer.

**A spacer** is a clear plastic tube. The inhaler fits in one end of the spacer and the other end of the spacer has a mouthpiece, or a mask which fits into the mouthpiece. A valve in the mouthpiece of the spacer opens and shuts as the person using it breathes in and out. The medicine goes directly into their lungs so that it can start working straight away.

Spacers greatly increase the effectiveness of the asthma medicine. You can get a spacer free of charge from your health provider.

## What if my child is very young or doesn't like the inhaler?


Children under the age of four can use their MDIs with a mask attached to the spacer. You or another adult will need to help them.

Once the spacer is in your child's mouth or the mask is over their mouth and nose, squirt the medicine into the spacer. It will remain suspended there, giving your child time to take six normal breaths through the mask.

The mask can come off the spacer when your child is able to use the spacer mouthpiece. For most children, this is by the age of four.

## Dry powder inhaler

The medicine in a **dry powder inhaler** is not a gas but a powder. This inhaler does not need to be shaken before you use it. The child breathes in the medicine directly from the inhaler's mouthpiece. Ask your doctor, nurse or pharmacist to show you how to use the inhaler.



## Using an inhaler with a spacer

The photographs below sets out six steps for using an MDI with a spacer. Younger children will need your help to follow these steps. As your child gets older, they will be able to do it themselves.



1 If using a mask, fit it to the spacer.



2 Hold the inhaler upright and give it a good shake.



3 Fit the inhaler into the opening at the end of the spacer.



4 Seal lips firmly around the mouthpiece or place the mask so it seals around the nose and mouth.



5 Press the inhaler once only.



6 Hold the spacer in place while taking six slow breaths through the mouth.

If your child needs another dose, repeat steps 2 to 6.

The medicine can only help your child if it gets into their lungs. It takes time to learn to use an inhaler, and people can forget. If you're not sure your child is using their inhaler correctly, ask your health provider to show you again.

### **How do I care for my child's spacer and inhaler?**

Wash the spacer once a week. This stops the medicine sticking to the side or blocking the valve. You should wash it in warm water with a mild detergent and then let it air dry.

Your child needs a new spacer every year. You can get this at the same time as you see your health provider about your child's action plan.

MDIs should also be washed weekly. You can use this time to check that there is still plenty of medicine left and that it hasn't reached the expiry date.

#### **Washing your MDI**

- 1** Remove the plastic mouthpiece cap.
- 2** Remove the metal canister but don't put it in the water.
- 3** Rinse the mouthpiece and cap under warm water for at least 30 seconds.
- 4** Shake off any excess water and allow to air dry thoroughly.
- 5** Put the metal canister back in and replace the cap.

## Introducing medicines



### What are the medicines for asthma?

There are several types of asthma medicine: preventer medicines, reliever medicines, combination medicines, and oral steroids. It's important to know the differences between these medicines, to take them correctly, and to be aware of the possible side effects. The tables on pages 16-19 will help you with this.

An 'oral steroid' is an asthma steroid medicine that is swallowed. It comes as a pill or a liquid.

Some children with asthma will be on a preventer medicine. Remember that asthma cannot be cured. This means it is important that your child keeps taking their preventer medicine as prescribed by your doctor.



Most asthma medicine is delivered through an inhaler. For this reason, people will often talk about 'using their reliever' or 'using their puffer' rather than 'taking their medicine'.

Preventer medicines contain steroids. Steroid medicines can have side effects, but these are mild and not nearly as bad as the effects of uncontrolled asthma. Talk to your health provider – your doctor, practice nurse, or asthma educator – if you're worried.

Even after asthma has been brought under control, it can flare up again. When that happens, you need to know which inhaler your child should use. Make sure you know the colour of your child's preventer inhaler and the colour of their reliever. You might want to write your own label to be sure.

Most children can manage their own asthma medicine from about the age of seven. However, even much older children need you to check that their asthma is under control and help watch for any changes.

Your child may only have been prescribed one or two types of medicine, or they may have more. Your doctor will write the names of your child's medicines on their asthma action plan. If you like, you could copy that information into the spaces in the table on the next page.

---

## **Preventer inhaler**

---

### **What it does**

A preventer is your child's most important medicine. It prevents the swelling and narrowing of the airways that causes asthma. This reduces the likelihood of an asthma attack.

Preventers work slowly, so your child won't notice any immediate change in how they feel. Over time, daily use will help to keep your child's asthma under control.

---

### **How to use it**

Preventers need to be taken every day or as prescribed, even when your child is well.

---

### **Possible side effects**

Possible side effects of preventer medicines include a sore throat, croaky voice, or a mouth infection. Your child can avoid these side effects by using a spacer and brushing their teeth or rinsing their mouth immediately after using their inhaler.

---

### **The preventer my child uses**



---

---

## Reliever inhaler

---

### What it does

A reliever medicine relaxes the tight bands of muscle around your child's airways to help air flow in and out more freely. This effect should last for about two to four hours.

### How to use it

Reliever medicines can be taken to relieve wheezing, coughing, or tightness in the chest area. Your health provider may also suggest that your child uses it before exercise.

Many people rely on their reliever inhalers to feel better immediately, but relievers do not treat the underlying cause of asthma symptoms. Uncontrolled asthma is serious and can even be life-threatening.

See your health provider if your child is using their reliever inhaler more than two or three times a week because this means their asthma is not under control.

### Possible side effects

Some side effects of reliever medicines include mild shaking, headaches, a racing heartbeat, and restlessness. If your child experiences these symptoms, see your doctor as soon as possible.

### The reliever my child uses



---

---

## Combination inhaler

---

### What it does

Combination inhalers contain two medicines so they help prevent asthma and relieve the symptoms.

---

### How to use it

Combination inhalers must be used every day, even when your child is well. Ask your health provider if this inhaler can be used in an emergency, because you may need a different reliever inhaler to use if your child is having an asthma attack.

---

### Possible side effects

The unwanted side effects of a combination inhaler can include mild shaking, headaches, a racing heartbeat, a sore throat, or an infection of the mouth. Your child can avoid getting a sore throat or fungal infection by using a spacer and brushing their teeth or rinsing their mouth immediately after taking their medicine.

---

### The combination inhaler my child uses



A large, empty light gray rectangular area intended for the user to write the name of the combination inhaler their child uses.

---

## Oral steroids

---

### What it does

Oral steroids (tablets and liquids) are very useful for bringing asthma symptoms under control when they are not settling with inhalers alone. They bring down the swelling in the lining of the child's airways and reduce the amount of mucus produced. Only some children with asthma need oral steroids.

### How to use it

Oral steroids are usually taken for about five days, along with your child's preventer medicine.


### Possible side effects

A short course of oral steroids is safe, with no lasting side effects. Some people notice a change in mood, energy levels, or appetite. These side effects are usually mild. You can discuss the possible short-term side effects and any other concerns with your doctor or pharmacist.

Very few children need long-term treatment with oral steroids. These children will be under the care of a specialist.

---

### The oral steroid my child uses



## What about complementary medicines?

It's OK to try alternative approaches, such as rongoā Māori or Chinese herbal medicine, provided you discuss them with your doctor and use them alongside and not instead of prescribed inhalers and medicines. Remedies containing natural products can be dangerous for people with asthma. Remedies like bee pollen or echinacea can be a trigger for asthma, and sometimes a natural remedy can interfere with your child's prescribed medicine.

On the other hand, traditional healing systems such as rongoā Māori are increasingly respected in mainstream medicine for taking a holistic view of health that includes the spirit, mind, body, and whānau. Some researchers are looking at the use of plant remedies, including those used by Māori, but so far there is little clear evidence about their effects. However, relaxation therapies, such as yoga, have been shown to help reduce asthma caused by stress.

If you're thinking about trying a natural remedy, find all the information you can and talk about it with your health provider. But remember – it's not safe to stop your child's medicine or give them a new medicine without first having that talk.

I love playing with my friends!  
They really help me.



---

# What should I do to manage my child's asthma?

Your doctor will have given you a child asthma action plan and you may also have been using a symptom diary. You can use the asthma action plan and symptom diary to help manage your child's asthma and prevent an attack.



The **action plan** tells you what to do when your child's asthma improves or gets worse. It tells you which medicine to use and how much. The amount and type of medicine will change with changes in your child's asthma symptoms. The table on page 23 shows how a child's symptoms may change from a mild asthma flare-up to an asthma emergency.

Your doctor will fill out your child's action plan, using the information you provide. This will help you to make the right decisions. Your doctor should update it at least once a year and any time there is a change in your child's medicine.

It is very important that you, your child, and all the people who care for your child know and understand the instructions in your child's action plan.



The **symptom diary** is a place where you can record your child's asthma symptoms so you can see any patterns over time. Seeing these patterns helps you and your health provider understand what is triggering your child's asthma and how well the medicine is working.

You don't need to fill in the symptom diary all the time.

However, it's very helpful when your child is first diagnosed, if there is a change in their medication, or if there seems to be a change in your child's symptoms. As your child gets older, you can help them take responsibility for their own symptom diary.



Both the asthma action plan and the symptom diary use a traffic light system. **Green** means that your child is good to go... the asthma is under control. **Orange** means that their asthma is getting worse and your child needs to start taking their reliever medicine. **Red** means that this is an emergency – your child is having an asthma attack and needs help.

	<p><b>Well</b></p> <p><b>When I'm well:</b></p> <ul style="list-style-type: none"> <li>I have no cough</li> <li>I play just like other children</li> <li>I use my reliever puffer less than 2 times a week</li> </ul>	<p><b>My puffers are:</b></p> <p><b>Preventer:</b> Take this every day even when I'm well.          The name of my preventer is _____ The colour is _____          I take _____ puffs in the morning and _____ puffs at night through a spacer.</p> <p><b>Reliever:</b> Take this only when I need it.          The name of my reliever is _____ The colour is _____          I take _____ puffs through a spacer when I wheeze, cough or when it's hard to breathe.</p> <p>If I find it hard to breathe when I exercise I should take _____ puffs of my reliever</p>
	<p><b>Worse</b></p> <p><b>When my asthma is getting worse:</b></p> <ul style="list-style-type: none"> <li>I cough or wheeze and it's hard to breathe, or</li> <li>I'm waking at night because of my asthma, or</li> <li>I cough or wheeze when I play or</li> <li>I cough or wheeze when I play or I use my reliever puffer to control my asthma more than 2 times per week</li> </ul>	<p><b>If my asthma gets worse I should:</b></p> <p>Keep taking my preventer every day as normal and take _____ puffs of my reliever every 4 hours          If I'm not getting better doing this I should see my doctor today</p> <p><b>Contact:</b></p> <div style="border: 1px solid black; height: 40px; width: 100%;"></div>
	<p><b>Worried</b></p> <p><b>My asthma is a worry when:</b></p> <ul style="list-style-type: none"> <li>My reliever isn't working, or</li> <li>I'm finding it hard to breathe, or</li> <li>I'm breathing hard and fast, or</li> <li>I'm tucking in around my ribs/throat, my looking under my skin</li> <li>I'm looking pale or blue</li> </ul>	<ul style="list-style-type: none"> <li>Sit me down and try to stay calm</li> <li>Give me 6 puffs of reliever through a spacer, taking 6 breaths for each puff</li> <li><b>If I don't start to improve I need help now</b></li> </ul>
<p><b>Emergency</b></p> <p><b>Do:</b> 111 and ask for an ambulance <b>WHILE YOU'RE WAITING</b></p> <ul style="list-style-type: none"> <li>Try to stay calm and keep me sitting upright</li> <li>Give 6 puffs of reliever through a spacer every 6 minutes with 6 breaths for each puff until help arrives</li> </ul>		
<p>Date Prepared: _____ Doctors Signature: _____ Plan to be reviewed when treatment changed</p>		

You can do all the right things, but your child may still have an asthma attack. Everything you are doing is helping but sometimes extra assistance is needed. In an emergency, help is at hand.

Please remember: If you have any questions, ask your doctor, nurse, or asthma educator. They're here to help!

---

# What should I do in an emergency?

In an emergency, try to stay calm. It helps to remember the six asthma steps: **A**ssess, **S**it, **T**reat, **H**elp, **M**onitor, and **A**ll OK!



## Assess

Assess whether the asthma attack is mild, moderate, or severe.

---

### Mild

Mild symptoms indicating that asthma is worsening can include:

- slight wheeze
- mild cough
- coughing or wheezing when excited or running

---

### Moderate

Moderate symptoms indicating that asthma is worrying can include:

- obvious breathing difficulties
- persistent cough
- difficulty speaking a complete sentence

---

### Severe

Severe symptoms indicating an asthma emergency can include:

- fear
- gasping for breath
- difficulty speaking more than one or two words
- looking pale and sounding quiet
- complaining that the reliever medicine is not working
- unresponsive – doesn't answer when spoken to

**Dial 111 for an ambulance!**



## Sit

Sit your child upright and stay with them. Lean them forward slightly and support their arms either on their knees or on a table.



You may like to cut this page out of the booklet and put it somewhere you can easily find it, such as on the fridge.



## Treat

Treat mild asthma symptoms with two puffs of a reliever inhaler. Treat moderate or severe symptoms with six puffs of the inhaler.

If the reliever medicine comes in a metered dose inhaler, use a spacer. Put the spacer into the child's mouth. Puff the inhaler once into the spacer and have the child breathe in and out six times.

Encourage your child to breathe as normally as possible.



## Help

If your child's asthma does not start to get better after six minutes, or if you or your child are frightened, call an ambulance. Continue giving your child six puffs of their reliever inhaler every six minutes until help arrives. Make sure they take six slow breaths for each puff. Keep doing this until they get better, or until the ambulance arrives.

Remember: **Six** puffs ... **six** breaths for each puff ... and repeat every **six** minutes.



## Monitor

Stay with your child and watch carefully, even if they seem to get better. If your child is not finding it easier to breathe, give repeat doses of the reliever inhaler and call an ambulance.



## All OK

Your child can return to quiet activities when they no longer wheeze, cough, or feel breathless. Keep monitoring their symptoms and take action if required, following the steps in the action plan. If your child's attacks are becoming more frequent or worrying, see your doctor as soon as possible.

---

## Should my child be physically active?



Exercise is good for us all, but did you know that it's especially good for people with asthma? Exercise can help improve the way their lungs work and boost their confidence so they can better manage their condition.

Unfortunately, exercise can trigger asthma, even in people whose asthma is well controlled. It's important that you, your child, and other responsible adults know what to do when this happens. These are some ways you can help to make exercise a safe part of your child's life:

- Talk with your health provider about your child's physical activity.
- If your child has been prescribed a dose of reliever medicine to take before exercise, check that they use it. The name of the reliever medicine and the dosage are written on your child's asthma action plan.
- Make sure your child takes their preventer medicine every day. Watch your child use their inhaler to check they are taking it correctly.
- Make sure your child always carries their reliever inhaler or has one nearby.
- If there is an adult in charge, tell them about your child's asthma action plan.
- Check that your child knows the signs of an asthma attack and when they should ask someone to dial 111 if you're not there.
- Encourage your child to choose sports and activities that have lots of stopping and starting, like tennis, softball, and most team sports.
- Encourage your child to warm up and warm down.
- Be aware that your child may need to avoid outside physical activity on cold, dry days or when there is a lot of pollen around.
- When it's cold, cover your child's nose and mouth with a scarf.



“

Alex Urlich<sup>2</sup> has a particularly severe form of asthma, but it doesn't stop him doing what he loves. Alex says:

For anyone who has asthma and is considering a sport, my advice to you is give it a go. If it's a strain, feel free to stop, but I promise you sports will make you strong and fit and feel like you can conquer anything!

”

<sup>2</sup> You can hear more from Alex by going to the digital learning resource located at [www.learnaboutlungs.org.nz](http://www.learnaboutlungs.org.nz)

---

## Who can help?

Your child is a precious taonga. We need all our children to grow up safe and strong. Keeping your child safe is a partnership. It is a partnership that you lead but that also includes your child, whānau, friends, health providers, and all the groups to which your child belongs, including their school or early childhood centre.

### Your child is Number 1!

When your child is very young, you need to take charge of their asthma management, but over time, your child needs to take control. Here are some things you can do to help:

- Asthma is a part of your child's life. Children's feelings about their asthma can be as important as the asthma itself. Encourage your child to talk about their thoughts, feelings, and experiences. Make sure your child knows they can still do most of the things other children can do. So long as they have a plan, they will be safe.
- The Asthma and Respiratory Foundation NZ has other resources available from their website and your health provider. Help your child to read and understand the information in these resources. Make sure they understand their own action plan.
- Teach your child the words they need for telling people when they think their asthma is getting worse. Try to include your child in conversations where you explain your child's asthma action plan to other people. Check that they know that if they feel they're getting worse, they can tell an adult who will help.



“

I taught Sione some simple things he could say to tell his teacher or other adults about how he was feeling. If I noticed his symptoms appearing, I'd say, "You look as if you're having trouble breathing. I can hear you making a whistling sound. That is called 'wheezing'. Does your chest feel tight? If you have a tight chest or you start to wheeze, please tell me or another adult. Let us know that you need your reliever."

By the time Sione started school, he knew to tell his teacher when he was 'wheezing' and needed his 'reliever'. I talked to the teacher and made sure she knew that my son knew what he was talking about and what she should do about it.

Helen, parent

”

## Involve your whānau and friends

Your whānau and friends all want to help their tamariki. Make sure they understand what asthma is and what they can do to help manage your child's asthma. Talk about your child's asthma action plan and put a copy somewhere they can see it. Get your child to show them how they use their reliever.

It will help your child if they know about people with asthma who are living active, healthy lives. Ask your whānau and friends to share stories that show your child that it can be done.

## Getting the best out of your health providers


In this booklet, we often talk about 'health providers'. These include doctors, nurses, and asthma educators. Together they form a team of people who are there to help you and your child. On pages 33–35, there is a directory you can complete so you can always remember the names of your child's health providers and how to contact them.

“ Sometimes I get asthma, but it's ka pai. We have a plan! ”



There is a lot to learn about asthma and how to deal with it. Your health provider is there to help you, but sometimes appointments can feel rushed or you can feel too anxious to take everything in. These are some things that can help:

- ✓ Before an appointment, write down your questions. Take a pen with you so you can jot down the answers.
- ✓ If you've been filling out a symptom diary, take it with you. Your health provider will use this to notice any patterns and think about how the asthma can be better managed.
- ✓ It's important to ask questions if you're unsure or there is something you don't understand. If an explanation isn't clear, ask your health provider to explain it another way. Maybe they could draw a picture or give you a booklet?
- ✓ Take a friend or whānau member so both of you can hear the advice and ask questions. The other person could also take notes.
- ✓ If possible, only take the child with asthma to the appointment and leave your other children with a caregiver. This will make it easier to concentrate.
- ✓ Keep a written record of your child's medical history along with information about your whānau. Does anyone else in your whānau have asthma? Take your written notes with you when you first see a new provider or if your child has to go to hospital.
- ✓ Once every year, your health provider is expected to look at your child's action plan and check that it is right. This is a good time to talk things over and for the provider to check that your child is using their inhaler correctly.
- ✓ Please don't feel bad if you need to ask for the same information again. There can be a lot to take in and it can be hard to remember, especially when you feel worried.



I was worried that Brooklyn wasn't using her inhaler properly. I got the nurse at the local medical centre to show us.

We talked over her asthma action plan and then before I left, she showed me how to use the inhaler again. She then asked me to do it with Brooklyn.

## Possible questions for your health provider

Some questions might include:

- 1 What is the medicine in my child's inhalers? What are the side effects? What can we do about the side effects?
- 2 What are some other things we can do to prevent my child's asthma attacks?
- 3 I think I can see some triggers for my child's asthma. How can we check? How can we avoid this trigger?



## Create an asthma-friendly community

Most children go to school or an early childhood centre and belong to other groups as well. It's important that the adults your child spends a lot of time with know about their asthma and what to do in an emergency. It's best if they read the action plan and talk it through with you.

We all want the children around us to be safe. All schools are auahi kore (smokefree), and many have asthma-friendly policies. This means they have chosen to make the school a safe and healthy place for people with asthma. Many marae, churches, and other organisations have also put in place asthma-friendly and auahi kore policies. If a group your child belongs to is interested in this, they could contact the Asthma and Respiratory Foundation NZ for information and resources.

---

# Who are our asthma health providers?

It helps to keep a written record of the people who provide health care to your child.

This directory has the names and contact details of some of the health providers who are part of your child's asthma team. In some places, there are blanks for you to fill in the names and contact details of people you see. The table doesn't include all possible health providers. The final row is blank so you can add other people who your child may see.

---

## Health provider

Your family **doctor** will be the first person you see to get a diagnosis and work out a plan for treating your child's asthma. The **practice nurse** can help you and your child to understand the plan and put it into action.

---

The **Asthma and Respiratory Foundation NZ** works to improve the respiratory health of all New Zealanders. You can find lots more information about asthma on the Foundation's website.



## Main contact

Names:

Address:

Phone number:

---

[www.asthmaandrespiratory.org.nz](http://www.asthmaandrespiratory.org.nz)

---

## Health provider

---

You can find an **asthma educator** by getting in touch with an asthma society or a trust. These services provide information and support, free or for a small charge. You can find them by asking your doctor or looking them up on the Asthma and Respiratory Foundation NZ website or in your phone book.

---

Your doctor may refer your child to a **specialist** with training in a particular area, such as treating respiratory diseases or diagnosing allergies or child health (a paediatrician).

---

Your doctor may refer your child to a **physiotherapist** to learn techniques to control breathing and remove mucus from their throat.



## Main contact

Names:

Address:

Phone number:

---

Names:

Address:

Phone number:

---

Names:

Address:

Phone number:

---

## Health provider

The Ministry of Health provides a directory you can use to find the contact details for **Māori health providers**.

---

Many services are available to support people who wish to quit smoking. **Quitline** is a great starting point.

---

Is there someone else your child sees about their asthma?  
You could write the details here.



## Main contact

[www.health.govt.nz/your-health/](http://www.health.govt.nz/your-health/)

Names:

Address:

Phone number:

---

[www.quit.org.nz](http://www.quit.org.nz)

Names:

Address:

Phone number:

---

Names:

Address:

Phone number:

Your child is surrounded by a team of people who care about their health. With your support and theirs, your child can manage their asthma and have a healthy, active life.





**Ehara taku toa i te toa  
takitahi, engari he toa  
takitini.**

**My strength is not  
mine alone, but that  
of many.**

**Asthma  
+ Respiratory**  
FOUNDATION NZ

Published by the Asthma and  
Respiratory Foundation NZ

**[asthmaandrespiratory.org.nz](http://asthmaandrespiratory.org.nz)**

All text and images copyright to the  
Asthma and Respiratory Foundation NZ.

All rights reserved. Enquiries should be  
made to the publisher.

SEPTEMBER 2017-ASTH59